

Unmetabolized Folic Acid, Tetrahydrofolate, and Colorectal Adenoma Risk

Judy R. Rees¹, Carolyn B. Morris², Janet L. Peacock^{1,3}, Per M. Ueland⁴, Elizabeth L. Barry¹, Gail E. McKeown-Eyssen⁵, Jane C. Figueiredo^{6,7}, Dale C. Snover⁸, and John A. Baron^{1,2}



Abstract

In a randomized trial of folic acid supplementation for the prevention of colorectal adenomas, we previously found indications of increased risk during later treatment and follow-up. This could have been due to the unmetabolized folic acid (UFA) or natural reduced and methylated folates (mF) to which it is metabolized. In *post hoc* analyses, we measured mF (the sum of 5-methyl-tetrahydrofolate and 4- α -hydroxy-5-methyl-THF) and UFA concentrations in the serum of 924 participants. Using binomial regression models with a log link, we assessed the associations between plasma mF or UFA and adenoma occurrence. We found no association between plasma mF or UFA and overall adenoma risk. However, during later follow-up, the pre-specified, composite endpoint of high-risk findings (advanced or multiple adenomas) was positively associated with plasma mF

($P_{\text{linear trend}} = 0.009$), with a 58% increased risk for participants in the upper versus lowest quartile. An irregular association was seen with plasma UFA, with suggestions of an inverse trend ($P_{\text{linear trend}} = 0.049$). A modest, significant inverse association was also seen between mF and risk of serrated lesions, with a 39% lower risk for upper versus lower quartile participants ($P_{\text{linear trend}} = 0.03$). In conclusion, during the later follow-up period in which folic acid supplementation was previously seen to increase the risk of advanced and multiple adenomas, higher serum mF was associated with a higher risk of multiple and/or advanced adenomas, but no clear indication that UFA played a direct role. There were indications that higher mF was associated with reduced risk of serrated polyps. *Cancer Prev Res*; 10(8): 451–8. ©2017 AACR.

Introduction

Extensive epidemiological and preclinical data have suggested that folates, a family of water-soluble vitamins that serve as cofactors for DNA synthesis and methylation reactions, exert substantial anti-neoplastic effects (1). Motivated by these findings, we conducted a randomized controlled trial to test the effect of 1 g/day folic acid for the prevention of colorectal adenoma recurrence (2). Although we found no effect during the first three years of treatment, we saw unexpected indications of increased risk during subsequent follow-up and treatment: the relative risk for the occurrence of advanced adenomas was 1.67 (95% CI,

1.00–2.80) and for 3 or more adenomas, 2.32 (95% CI, 1.23–4.35; ref. 2).

Although our finding of increased adenoma risk from folic acid supplementation may have been due to chance, there are other indications of increased cancer-related risks of folic acid, depending on dose, timing, and folate status before supplementation. Whereas moderate doses may reduce risk, carcinogenesis may be enhanced by larger doses, or by supplementation after a neoplastic lesion has developed (3–6). The mechanisms for this adverse effect have not been fully elucidated but may include the provision of nucleotides for carcinogenesis (7).

Folic acid is a stable, oxidized molecule rarely found in nature, though widely used for food fortification and in supplements. Folic acid is the molecular backbone of folates, but it is inactive until it is metabolized into the natural reduced forms, including 5-methyltetrahydrofolate (5m-THF), the prevailing circulating folate species (8). Because folic acid enters the folate metabolic cycles differently than 5m-THF and has a distinct metabolism, it may have biological effects that differ from those of the naturally occurring folates. For example, some studies suggest that UFA reduces natural killer cell cytotoxicity *in vitro* (9, 10). Thus, the increased risks associated with folic acid supplementation in our study could have been the result of folic acid itself, not the 5m-THF generated by it.

For both 5m-THF and folic acid, genetic variants in folate metabolism may also be pertinent. For example, the homozygous C677T variant of methylenetetrahydrofolate reductase (MTHFR), which metabolizes methylenetetrahydrofolate to mTHF, may modify the association of folate status with risk of colorectal carcinoma and adenoma (11–13). MTRR, methionine synthase

¹Department of Epidemiology, Geisel School of Medicine at Dartmouth, Dartmouth-Hitchcock Medical Center, Lebanon, New Hampshire. ²Department of Medicine, University of North Carolina School of Medicine, Chapel Hill, North Carolina. ³Division of Health and Social Care Research, King's College London, London, United Kingdom. ⁴Section for Pharmacology, University of Bergen, Hordaland, Norway. ⁵Dalla Lana School of Public Health, University of Toronto, Toronto, Canada. ⁶Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California. ⁷Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, California. ⁸Department of Pathology, Fairview Southdale Hospital, Edina, Minnesota.

Note: Supplementary data for this article are available at Cancer Prevention Research Online (<http://cancerprevres.aacrjournals.org/>).

Corresponding Author: Judy R. Rees, Geisel School of Medicine at Dartmouth College, HB 7927, 1 Medical Center Drive, Lebanon, NH 03756. Phone: 603-653-3683; E-mail: Judy.Rees@dartmouth.edu

doi: 10.1158/1940-6207.CAPR-16-0278

©2017 American Association for Cancer Research.

reductase, maintains methionine synthase in its active form, and in this state maintains intracellular folate levels (14).

It is widely recognized that colorectal cancer is a heterogeneous disease, with two distinct precursor pathways: the conventional pathway (accounting for about 60% of colorectal cancers) and the serrated pathway, which leads to CpG island methylated colorectal cancers (accounting for the remaining ~40%; refs. 15, 16). Because these involve distinct genetic and epigenetic processes, it is possible that folates affect these pathways in different ways.

In a secondary, observational analysis, we previously found no consistent association between folate status and adenoma risk (17). However, during early follow-up, total folate intake and plasma total folate were inversely associated with adenoma risk among subjects randomized to folic acid placebo, while in later follow-up there were inconsistent suggestions of increased risks for dietary folate intake and red blood cell folate. In this analysis, we extend our analyses using separate measures of plasma UFA and methylated folate (mF), and separate assessment of the conventional adenoma pathway and the serrated pathway. We hypothesized that long-term risk after folic acid supplementation is independently associated with plasma UFA levels.

Materials and Methods

Briefly, we conducted a double-blinded, placebo-controlled, randomized trial (NCT00272324) in a 3×2 factorial design to compare the effects of oral aspirin (81 mg/day, 325 mg/day, or placebo) with placebo or folic acid 1 g/day, in the prevention of colorectal adenomas in persons with a history of adenomas (2, 18). The trial's primary outcome was the occurrence of ≥ 1 colorectal adenomas. Secondary outcomes were the occurrence of advanced adenomas (tubulovillous or villous adenomas, or adenomas that were large (≥ 1 cm) or had invasive cancer or high-grade dysplasia); high-risk findings (≥ 1 advanced adenomas or ≥ 3 adenomas); and serrated lesions (comprising hyperplastic polyps, sessile serrated adenomas/polyps, and traditional serrated adenomas). The recruitment period (1994–98) coincided with the beginning of folic acid food fortification in the United States and Canada that became mandatory in 1998 (19, 20). All participants provided written informed consent; the trial was approved by each participating center's institutional review board.

Participants aged 21–80 with a recent adenoma and no known remaining adenomas after a complete colonoscopy were identified at nine clinical centers, as described previously (18). Exclusion criteria included vitamin B₁₂ deficiency, contraindication to aspirin treatment, or ongoing treatment with aspirin or related compounds (2, 18).

After a run-in phase of approximately 3 months, 1,021 participants were randomized to aspirin (81 mg, 325 mg or placebo) and folic acid (1 mg or placebo); 100 early participants were randomized only to aspirin/placebo. The initially planned study interventions continued until a follow-up colonoscopy anticipated approximately 3 years after the qualifying examination. This demarcated the end of the first follow-up interval and the beginning of the second interval, which lasted until 6 years later, one year longer than a 5-year colonoscopic surveillance cycle (Fig. 1).

At the end of the first follow-up interval, participants were asked to continue placebo/folic acid treatment until their next surveillance colonoscopy, anticipated 3 to 5 years later; 729/1,084

(67%) agreed to do so (Fig. 1). Of the 355 who chose to discontinue randomized treatment, 278 (78%) agreed to observational follow-up. Participants still on study treatment were asked to stop taking study tablets by October 1, 2004, because folic acid supplementation had been associated with increased adenoma risk.

Plasma measures

Blood samples were collected at enrollment ("year 0," reflecting prerandomization folate status), and shortly before the end of the first treatment interval ("year 3"), reflecting folate status on treatment. The blood draws were not timed to intake of food or study tablets. This study examined associations with circulating concentrations of folic acid and naturally occurring folates shortly before the year 3 colonoscopy.

Plasma levels of 5m-THF, 4- α -hydroxy 5-methyl-THF (hm-THF), and unmetabolized folic acid (UFA) were analyzed at Bevilal (21), using LC MS/MS (22). hm-THF is formed upon degradation of 5m-THF during storage so we pooled these measures for each sample and refer to the sum as mF (22). The levels of detection were 0.27 nmol/L for UFA, 0.20 nmol/L for hm-THF, and 0.07 nmol/L for 5m-THF (22). The intra-run CVs ranged from 5.0% to 9.9%, and the inter-run CVs from 3.3% to 9.5% (22). We identified the 677C>T, V222A (rs1801133) polymorphism of the methylenetetrahydrofolate reductase (*MTHFR*) gene and the 66A>G, I22M (rs1801394) polymorphism of methionine synthase reductase (*MTRR*) as previously described (23).

Follow-up


Participants taking study treatments were contacted every 4 months for information on adherence, use of personal nutritional supplements including folic acid, large bowel endoscopies and major medical events. During observational follow-up, they were contacted annually. For all tissue removed from the bowel, slides were obtained if possible and sent to a single, blinded pathologist for classification.

Statistical analysis

We used conventional descriptive statistics to summarize characteristics of the study population. Correlations were assessed using Kendall's Tau b. We used the circulating folate measures at year 3 to characterize the folic acid and folate status of subjects during the first follow-up interval. In the analysis for the second follow-up interval, we included participants whose study folic acid treatment remained unchanged from that in the first interval: those randomized to folic acid who agreed to continue study treatment, those allocated folic acid placebo whether or not they agreed to continue study pills, and 100 participants randomized only to aspirin or placebo (i.e., no randomization to folic acid/placebo). Thus, apart from changes in diet, and non-compliance with the study treatment regimen, the year 3 blood specimen represented the UFA and mF status during the second follow-up interval as well as the first.

We fit binomial regression models with a log link to describe the associations of circulating folate measures at year 3 with the occurrence of adenomas and serrated polyps during each of the surveillance intervals. The endpoints were ≥ 1 adenomas; ≥ 1 advanced adenomas; high-risk findings; ≥ 1 serrated lesions. All analyses were adjusted for age and sex, but not for treatment group (which would be reflected in the year 3 plasma measurements).

Figure 1.
Design of the trial and participants studied in this analysis (adapted from Cole et al., ref. 2).



Clinical center did not materially affect the analyses, and was excluded from all models. Personal folic acid supplement use during the second follow-up interval did not contribute usefully to the models for any of the measured outcomes, either as a covariate or as an effect modifier. It was therefore omitted from the analysis, and the same model was used for both intervals (i.e., age, sex, mF, and UFA). Estimates for mF and UFA were mutually adjusted for each other so that the risk ratios reflected the independent associations of each analyte. mF was modeled in quartiles; UFA was modeled in four categories, undetectable (set to zero) plus tertiles of the remaining values. We assessed patterns of associations with a general test of heterogeneity (Wald test) based on the categorical variables. We assessed tests for trend using the linear and quadratic functions of the continuous measurements. This approach allowed testing of both monotonic trends and the possibility that modest increases in mF or UFA levels might be beneficial, while higher levels detrimental. Results are given as relative risks of outcome by mF or UFA category. The possibility of effect modification by randomized aspirin assignment (dichotomous) or by *MTHFR* or *MTRR* polymorphisms was assessed using likelihood ratio tests. All analyses were performed with SAS v9.4.

Results

Year 3 plasma measurements and colonoscopy data were available for 924 participants in the first interval, 680 of whom completed a further 3 to 6 years of colonoscopic follow-up (Fig. 1; Table 1). At enrollment, the mean concentration of mF was 37.7 nmol/L (SD 22.4); at year 3 this had increased to 46.3 nmol/L (SD 21.7) in the placebo group and 84.8 nmol/L (SD 29.5) in the folic acid treatment group. At enrollment, UFA was detected in 19.3% of participants, with a mean concentration

of 4.2 (6.5) nmol/L for the 170 with detectable levels. Just before the 3-year surveillance colonoscopy, detectable UFA was found in 23.2% of the placebo group and 73.0% of those in the folic acid group, with mean concentrations of 3.4 (8.5) and 21.7 (24.9) nmol/L for those with detectable levels, respectively. At year 3, mean plasma mF increased from 48.0 to 97.0 nmol/L within increasing categories of UFA (Supplementary Fig. S1). The correlation between mF and UFA at year 3 was $r = 0.46$ ($P < 0.0001$).

There were no significant associations between circulating UFA or mF levels and overall adenoma risk during either follow-up interval (Table 2). However, during the second follow-up interval, there were indications of an association of mF with increasing risk of advanced adenomas and high risk findings (advanced or ≥ 3 adenomas; Table 2). For high-risk findings, there was a linear trend of increasing risk with increasing levels, with a 58% increased risk for participants in the upper quartile compared with the lowest (Table 2). There was a weaker, borderline-significant trend for advanced adenomas (25% increased risk for upper vs. lowest quartile). During the second interval, UFA showed an erratic association with high-risk findings, with a borderline-significant inverse linear trend ($P = 0.049$). Analyses without mutual adjustment for mF and UFA showed broadly similar findings, though with nonsignificant trends (Supplementary Data; Table 1).

There was no evidence for any differences between men and women in the association of mF or UFA with any of these adenoma endpoints (all $P_{\text{interaction}} > 0.15$; Supplementary Data; Table 2). There was also little evidence for a quadratic relationship between mF or UFA and these outcomes during early or later follow-up (all $P > 0.15$; not shown). No significant interactions were seen for any outcome between randomized aspirin use (dichotomous) and either UFA or mF.

Rees et al.

Table 1. Characteristics of trial participants included in serum analyses

Enrollment	Participants who completed interval 1 ^a (N = 924)		Participants who completed interval 2 ^a (N = 680)	
	Mean (SD)	N (%)	Mean (SD)	N (%)
Age	57.5 (9.3)		57.6 (9.0)	
Sex:				
Male		587 (63.5)		440 (64.7)
Female		337 (36.5)		240 (35.3)
Race/ethnicity				
Non-Hispanic white		784 (84.8)		591 (86.9)
Non-Hispanic black		54 (5.8)		33 (4.9)
Hispanic		56 (6.1)		35 (5.2)
Asian/Pacific Islander		22 (2.4)		15 (2.2)
Native American		4 (0.4)		3 (0.4)
Other		4 (0.4)		3 (0.4)
Year 0		N = 879		N = 649
mF ^b mean (SD) nmol/L	37.7 (22.4)		37.7 (22.4)	
Participants with detectable UFA ^c (%)		170 (19.3%)		119 (18.3%)
UFA mean (SD) for those with detectable levels, nmol/L	4.2 (6.5)		4.5 (7.3)	
Year 3		N = 924		N = 680
mF mean (SD), nmol/L	64.1 (32.0)		63.7 (31.7)	
Folic acid	84.8 (29.5)		87.0 (27.9)	
Placebo	46.3 (21.7)		46.3 (21.7)	
Number of participants with detectable UFA (%)		428 (46.3%)		314 (46.2%)
Folic acid		313 (73.0%)		230 (79.4%)
Placebo		115 (23.2%)		84 (21.5%)
UFA mean (SD) for those with detectable levels, nmol/L	16.8 (23.2)		16.7 (22.9)	
Folic acid	21.7 (24.9)		21.4 (24.5)	
Placebo	3.4 (8.5)		3.8 (9.2)	

^aInterval 1 is year 0 through the year 3 colonoscopy. Interval 2 began after the year 3 colonoscopy and ended 6 years later.

^bmF is methylated folate, the sum of measured 5-methyl-THF and 4- α -hydroxy 5-methyl-THF.

^cUFA is unmetabolized folic acid.

Associations for serrated pathway lesions differed markedly from those for adenomas. During the second surveillance interval, the risk of any serrated lesion was significantly inversely associated with plasma mF, with a 39% lower risk for upper vs. lower quartile participants (Table 2). There was no association with plasma UFA (Table 2).

As reported previously for the entire study population (24), there were no significant associations between adenoma risk and polymorphisms of *MTHFR* or *MTRR* in either follow-up interval (Supplementary Table S3). There were also no indications of interactions between plasma levels of mF or UFA and genetic variants in either gene with regard to risk of all adenomas (Supplementary Table S3). However, in the corresponding analysis of high risk-findings, there was an interaction of *MTHFR* with mF ($P = 0.03$) during the first follow-up interval, with higher mF-associated risk in CC participants than in those with the CT/TT genotype. (Supplementary Table S4). In the second interval, there was a broadly similar pattern, but the interaction did not reach statistical significance ($P = 0.08$).

Discussion

In our clinical trial, we previously found that participants randomized to 1 mg/day folic acid had an increased risk of advanced and multiple adenomas during longer follow-up (2). In this post hoc analysis, high circulating levels of mF were positively associated with the composite high-risk endpoint of advanced and/or multiple adenomas during this same period, while there was lower risk in the highest category of UFA, with a borderline significant trend. These findings suggest that the increased risks seen after folic acid supplementation (2) were due to the high mF levels, rather than due to a direct effect of

UFA itself (Fig. 2). In contrast, our findings for serrated lesions suggested a protective effect of higher levels of mF during later follow-up.

We found an interaction between *MTHFR* 677C→T and mF levels such that in the first follow-up interval there was a particularly elevated risk of high-risk findings associated with high mF only among individuals homozygous for the CC allele. This is consistent with our finding from the second follow-up interval that mF is associated with increased risk, as the enzyme variant associated with the T allele generates 5-methyltetrahydrofolate from 5, 10-methylenetetrahydrofolate less efficiently than the variant associated with the C allele. However, during the second follow-up interval, the interaction was less marked.

Cho and colleagues identified no overall association between circulating UFA and colorectal cancer, but with significant heterogeneity by sex, with higher risks among men and lower risks among women, with increasing UFA (25). In the second follow-up interval of our study, the directions of effects were similar to those seen by Cho and colleagues, but the interactions were not statistically significant in our smaller study population. Cho and colleagues also reported a significant interaction between UFA and *MTHFR* C677T, with a colorectal cancer OR of 2.20 (95% CI, 1.22–3.94) for the highest versus lowest levels of UFA among those with CT or TT genotype. Increasing risks associated with the TT or CT genotype have been found previously, either as main effects, or with increasing total folate levels (11, 26). We also found only slightly and nonsignificantly higher risk ratios in those with these genotypes at highest UFA levels.

Several laboratory studies indicate that folate or folic acid may enhance carcinogenesis in some settings. For example, in mouse

Table 2. Associations of plasma mF and UFA levels with risk of adenomas and serrated lesions

All adenomas ^b	Interval 1 ^a		Interval 2 ^a	
	N (events)	RR ^c (95% CI)	N (events)	RR ^c (95% CI)
mF nmol/L				
0-<37.35	218 (86)	1.00 (ref)	170 (62)	1.00 (ref)
37.35-<60.44	224 (91)	1.02 (0.79-1.33)	157 (59)	1.00 (0.73-1.39)
60.44-<85.50	221 (91)	1.04 (0.78-1.38)	167 (68)	1.06 (0.75-1.50)
≥85.50	224 (87)	1.02 (0.74-1.42)	162 (60)	1.01 (0.68-1.50)
		$P_{\text{heterogeneity}} = 0.99^{\text{d}}$		$P_{\text{heterogeneity}} = 0.98^{\text{d}}$
		$P_{\text{linear trend}} = 0.70$		$P_{\text{linear trend}} = 0.69$
UFA nmol/L				
0	475 (190)	1.00 (ref)	353 (130)	1.00 (ref)
>0-<3	163 (69)	1.04 (0.81-1.33)	111 (38)	0.88 (0.62-1.25)
3-<20	120 (51)	1.00 (0.74-1.35)	91 (45)	1.31 (0.95-1.82)
≥20	129 (45)	0.83 (0.58-1.18)	101 (36)	0.96 (0.64-1.43)
		$P_{\text{heterogeneity}} = 0.64^{\text{d}}$		$P_{\text{heterogeneity}} = 0.14^{\text{d}}$
		$P_{\text{linear trend}} = 0.13$		$P_{\text{linear trend}} = 0.81$
High-risk findings^e				
mF nmol/L				
0-<37.35	214 (25)	1.00 (ref)	170 (15)	1.00 (ref)
37.35-<60.44	222 (35)	1.35 (0.87-2.11)	157 (19)	1.13 (0.71-2.36)
60.44-<85.50	216 (26)	1.02 (0.61-1.71)	167 (25)	1.56 (0.84-2.90)
≥85.50	223 (32)	1.32 (0.77-2.24)	162 (21)	1.58 (0.80-3.09)
		$P_{\text{heterogeneity}} = 0.38^{\text{d}}$		$P_{\text{heterogeneity}} = 0.52^{\text{d}}$
		$P_{\text{linear trend}} = 0.96$		$P_{\text{linear trend}} = 0.009$
UFA nmol/L				
0	467 (63)	1.00 (ref)	353 (38)	1.00 (ref)
>0-<3	160 (21)	0.94 (0.60-1.46)	111 (16)	1.11 (0.65-1.89)
3-<20	119 (20)	1.17 (0.73-1.88)	91 (18)	1.46 (0.86-2.50)
≥20	129 (14)	0.73 (0.41-1.29)	101 (8)	0.59 (0.28-1.24)
		$P_{\text{heterogeneity}} = 0.45^{\text{d}}$		$P_{\text{heterogeneity}} = 0.10^{\text{d}}$
		$P_{\text{linear trend}} = 0.46$		$P_{\text{linear trend}} = 0.049$
Advanced adenoma^f				
mF nmol/L				
0-<37.35	214 (19)	1.00 (ref)	166 (13)	1.00 (ref)
37.35-<60.44	221 (24)	1.28 (0.78-2.10)	150 (14)	1.11 (0.59-2.07)
60.44-<85.50	214 (18)	1.09 (0.62-1.91)	162 (17)	1.18 (0.61-2.28)
≥85.50	222 (23)	1.56 (0.88-2.77)	158 (15)	1.25 (0.61-2.55)
		$P_{\text{heterogeneity}} = 0.39^{\text{d}}$		$P_{\text{heterogeneity}} = 0.94^{\text{d}}$
		$P_{\text{linear trend}} = 0.73$		$P_{\text{linear trend}} = 0.051$
UFA nmol/L				
0	466 (48)	1.00 (ref)	341 (28)	1.00 (ref)
>0-<3	159 (16)	0.91 (0.56-1.47)	108 (13)	1.35 (0.76-2.38)
3-<20	119 (11)	0.80 (0.44-1.41)	89 (13)	1.60 (0.88-2.94)
≥20	127 (9)	0.60 (0.29-1.09)	98 (5)	0.55 (0.23-1.32)
		$P_{\text{heterogeneity}} = 0.40^{\text{d}}$		$P_{\text{heterogeneity}} = 0.07^{\text{d}}$
		$P_{\text{linear trend}} = 0.24$		$P_{\text{linear trend}} = 0.07$
Serrated lesions^g				
mF nmol/L				
0-<37.35	223 (74)	1.00 (ref)	167 (52)	1.00 (ref)
37.35-<60.44	222 (47)	0.63 (0.44-0.90)	155 (43)	0.88 (0.60-1.28)
60.44-<85.50	219 (66)	0.87 (0.61-1.25)	163 (44)	0.90 (0.60-1.37)
≥85.50	226 (66)	0.85 (0.57-1.26)	158 (28)	0.61 (0.36-1.23)
		$P_{\text{heterogeneity}} = 0.09^{\text{d}}$		$P_{\text{heterogeneity}} = 0.30^{\text{d}}$
		$P_{\text{linear trend}} = 0.66$		$P_{\text{linear trend}} = 0.03$
UFA nmol/L				
0	478 (134)	1.00 (ref)	346 (98)	1.00 (ref)
>0-<3	162 (45)	1.03 (0.72-1.45)	112 (34)	1.18 (0.81-1.73)
3-<20	121 (38)	1.12 (0.76-1.66)	87 (17)	0.78 (0.46-1.35)
≥20	129 (36)	0.98 (0.64-1.50)	98 (18)	0.82 (0.47-1.45)
		$P_{\text{heterogeneity}} = 0.93^{\text{d}}$		$P_{\text{heterogeneity}} = 0.43^{\text{d}}$
		$P_{\text{linear trend}} = 0.66$		$P_{\text{linear trend}} = 0.91$

^aInterval 1 is year 0 through the year 3 colonoscopy. Interval 2 began after the year 3 colonoscopy and ended 6 years later.

^bConventional adenomas: tubulovillous, villous or tubular adenomas (i.e., non-serrated).

^cRisk ratios are estimated for each quartile of mF (or UFA), with respect to the first quartile as referent, in the following model: colonoscopy findings = age + sex + quartiles of mF (or UFA) + continuous UFA (or mF).

^d $P_{\text{heterogeneity}}$ is obtained from the Wald test based on categorical variables; $P_{\text{linear trend}}$ is based on linear function of the continuous data.

^eHigh-risk findings: 3 or more adenomas or advanced adenoma.

^fAdvanced adenoma: tubulovillous or villous adenomas, or adenomas that were large (≥1 cm), or had invasive cancer or high-grade dysplasia.

^gSerrated lesions: hyperplastic polyps, sessile serrated adenomas/polyps, and traditional serrated adenomas.

Rees et al.




Figure 2.
Possible causal pathways under investigation.

models of colorectal cancer, low doses of dietary folic acid were protective if given before tumors developed, but promoted further tumor growth if given after tumors had emerged (27), and in a rat model of breast cancer, folate deficiency was protective (28, 29). Other *in vitro* studies have shown that folate repletion after deficiency has the potential to lead to carcinogenesis (5). Some argue that, because folate antagonists are effective anticancer drugs, the rationale for folate supplementation to prevent cancer may be flawed; indeed, the use of antifolates in leukemia was first prompted by the observation that large doses of folic acid made patients with leukemia substantially worse (30).

Although the inverse association between high-risk findings and UFA may be a chance finding, there are reasons to speculate that synthetic folic acid might have different biological effects than naturally occurring folates (31, 32), some of them potentially harmful (33). UFA has been associated with neurological and cognitive effects in individuals with lower B₁₂ status (34, 35) and with reduced natural killer cell cytotoxicity in some studies *in vivo* (9, 10), although not *in vitro* (36). There are differences in the carrier systems through which these natural folates and UFA enter the cell, the points at which they enter the folate cycle, and the bioavailability of folates from diet and folic acid supplements (31, 37). The assimilation of folic acid into tetrahydrofolate is catalyzed by dihydrofolate reductase (DHFR), whose activity varies considerably between individuals and is inhibited by UFA (38). In addition, UFA may inhibit MTHFR, the enzyme that produces methyltetrahydrofolate (39). Thus, it is possible that UFA has tissue effects that decrease the bioavailability of intracellular mF levels following the supplementation. Further, after administration of radiolabeled folic acid to 10 volunteers, 40% of the increase in plasma total folate was due to mobilization of endogenous (unlabeled) folate rather than conversion of the ingested folic acid (40). The physiological differences between UFA and naturally occurring folate species are incompletely understood and require further study. The irregular risk ratios seen in the second interval of our study may also merit further investigation in future studies, to assess the possibility of a threshold effect.

After folic acid supplementation, 73.0% of our participants had detectable UFA, compared with 23.2% of controls; these figures are consistent with the wide range of detectable UFA seen in other studies (41–44). UFA has been found in the serum following boluses of dietary folic acid as low as 200 mcg in folate-replete women (34), suggesting that it should be expected in the serum of individuals taking 1 mg folate/day (45). A variety of factors influence the concentration of mF and UFA, including the time since ingestion of the supplement (46–48), variability in dietary intake including food fortification and DHFR polymorphisms. We did not control the timing of blood tests relative to pill-taking

or meals in our study; variable timing would increase the variability of plasma measures and reduce our ability to identify associations if any exist. This would be particularly true for UFA, which has a short half-life (~1 hour; ref. 49) as compared with the more deep folate compartments that have previously been detected after chronic supplementation with deuterium labeled folic acid (50). A future study should time blood draws consistently in relation to intake of folic acid from foods and supplements.

Our approach had the advantage of adjustment for serum mF while analyzing the effect of UFA and vice versa. By summing the measured concentrations of hm-THF and 5m-THF, we minimized the measurement error associated with 5m-THF breakdown to hm-THF in stored samples, (22). We also used a single blood sample during intervention (near the end of the first follow-up interval) to represent UFA or mF levels during two observation cycles; this is more reflective of in-trial levels than the presupplementation samples used in our previous report (17). However, a single measurement is an imperfect correlate of long-term exposure and may cause regression dilution bias. During a multiyear study, changes in adherence and in diet would likely reduce the reliability of the single blood measure in representing long-term folate status, and this would tend to reduce our ability to identify associations (51). A further limitation is that our findings with respect to adenoma and especially serrated polyps apply only indirectly to the study of colorectal cancer. Finally, in exploring trends in these data, we performed many significance tests, with the possibility of type 1 errors, particularly for the quadratic trend tests that were conducted; one of 20 tests was statistically significant with $P = 0.04$.

Conclusion

Our results indicate that the increased risk of advanced/multiple adenomas in later follow-up of our trial was likely related to higher levels of plasma mF, and not due to UFA. In contrast, findings regarding serrated lesions provide some suggestions of protection from mF in the serrated pathway. While our data do not indicate any significant harm associated with UFA itself, the associated high levels of mF seem to be detrimental with respect to adenoma risk. The widespread use of folic supplementation and food fortification makes associations between the levels of folate species and disease an important avenue for further research.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: J.R. Rees, J.A. Baron

Development of methodology: J.R. Rees, P.M. Ueland

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): G.E. McKeown-Eyssen, J.C. Figueiredo, J.A. Baron

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): J.R. Rees, C.B. Morris, J.L. Peacock, G.E. McKeown-Eyssen, J.C. Figueiredo, J.A. Baron

Writing, review, and/or revision of the manuscript: J.R. Rees, C.B. Morris, J.L. Peacock, P.M. Ueland, E.L. Barry, G.E. McKeown-Eyssen, J.C. Figueiredo, D.C. Snover, J.A. Baron

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): D.C. Snover

Study supervision: J.A. Baron

Acknowledgments

The authors wish to acknowledge the participants and staff of the Polyp Prevention Study.

Grant Support

This work was supported in part by grants 5 R01CA059005 and U54 CA100971 from the NIH. Study tablets were provided by Wyeth Consumer Health Care, Madison, NJ. J.L. Peacock is supported by the NIHR Biomedical

Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London, UK.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received October 31, 2016; revised April 24, 2017; accepted June 5, 2017; published OnlineFirst June 9, 2017.

References

- Giovannucci E. Epidemiologic studies of folate and colorectal neoplasia: a review. *J Nutr* 2002;132:2350S–5S.
- Cole BF, Baron JA, Sandler RS, Haile RW, Ahnen DJ, Bresalier RS, et al. Folic acid for the prevention of colorectal adenomas: a randomized clinical trial. *JAMA* 2007;297:2351–9.
- Song J, Medline A, Mason JB, Gallinger S, Kim YI. Effects of dietary folate on intestinal tumorigenesis in the *apcMin* mouse. *Cancer Res* 2000;60:5434–40.
- Song J, Sohn KJ, Medline A, Ash C, Gallinger S, Kim YI. Chemopreventive effects of dietary folate on intestinal polyps in *Apc+/-Msh2-/-* mice. *Cancer Res* 2000;60:3191–9.
- Melnik S, Pogribna M, Miller BJ, Basnakian AG, Pogribny IP, James SJ. Uracil misincorporation, DNA strand breaks, and gene amplification are associated with tumorigenic cell transformation in folate deficient/repleted Chinese hamster ovary cells. *Cancer Lett* 1999;146:35–44.
- Folate Kim YI. colorectal carcinogenesis, and DNA methylation: lessons from animal studies. *Environ Mol Mutagen* 2004;44:10–25.
- Choi SW, Mason JB. Folate and carcinogenesis: an integrated scheme. *J Nutr* 2000;130:129–32.
- Bailey LB, Stover PJ, McNulty H, Fenech MF, Gregory JF, Mills JL 3rd, et al. Biomarkers of nutrition for development-folate review. *J Nutr* 2015;145:1636S–80S.
- Troen AM, Mitchell B, Sorensen B, Wener MH, Johnston A, Wood B, et al. Unmetabolized folic acid in plasma is associated with reduced natural killer cell cytotoxicity among postmenopausal women. *J Nutr* 2006;136:189–94.
- Sawaengsri H, Wang J, Reginaldo C, Steluti J, Wu D, Meydani SN, et al. High folic acid intake reduces natural killer cell cytotoxicity in aged mice. *J Nutr Biochem* 2016;30:102–7.
- Chen J, Giovannucci E, Kelsey K, Rimm EB, Stampfer MJ, Colditz GA, et al. A methylenetetrahydrofolate reductase polymorphism and the risk of colorectal cancer. *Cancer Res* 1996;56:4862–4.
- Ulrich CM, Kampman E, Bigler J, Schwartz SM, Chen C, Bostick R, et al. Colorectal adenomas and the C677T MTHFR polymorphism: evidence for gene-environment interaction? *Cancer Epidemiol Biomarkers Prev* 1999;8:659–68.
- Kim DH. The interactive effect of methyl-group diet and polymorphism of methylenetetrahydrofolate reductase on the risk of colorectal cancer. *Mutat Res* 2007;622:14–8.
- Leclerc D, Wilson A, Dumas R, Gafuik C, Song D, Watkins D, et al. Cloning and mapping of a cDNA for methionine synthase reductase, a flavoprotein defective in patients with homocystinuria. *Proc Natl Acad Sci U S A* 1998;95:3059–64.
- Fearon ER, Vogelstein B. A genetic model for colorectal tumorigenesis. *Cell* 1990;61:759–67.
- Jass JR. Classification of colorectal cancer based on correlation of clinical, morphological and molecular features. *Histopathology* 2007;50:113–30.
- Figueiredo JC, Levine AJ, Grau MV, Barry EL, Ueland PM, Ahnen DJ, et al. Colorectal adenomas in a randomized folate trial: the role of baseline dietary and circulating folate levels. *Cancer Epidemiol Biomarkers Prev* 2008;17:2625–31.
- Baron JA, Cole BF, Sandler RS, Haile RW, Ahnen D, Bresalier R, et al. A randomized trial of aspirin to prevent colorectal adenomas. *N Engl J Med* 2003;348:891–9.
- US Department of Health and Human Services Food and Drug Administration. Food standards: amendment of the standards of identity for enriched grain product to require addition of folic acid. *Fed Regist* 1996;61:8781.
- Regulations Amending the Food and Drug Regulations (1066). *Canada Gazette Part II* 1998;132:SOR/98–550.
- Mason JB, Choi SW. Folate and carcinogenesis: developing a unifying hypothesis. *Adv Enzyme Regul* 2000;40:127–41.
- Hannisdal R, Ueland PM, Eussen SJ, Svardal A, Hustad S. Analytical recovery of folate degradation products formed in human serum and plasma at room temperature. *J Nutr* 2009;139:1415–8.
- Figueiredo JC, Levine AJ, Grau MV, Midttun O, Ueland PM, Ahnen DJ, et al. Vitamins B2, B6, and B12 and risk of new colorectal adenomas in a randomized trial of aspirin use and folic acid supplementation. *Cancer Epidemiol Biomarkers Prev* 2008;17:2136–45.
- Levine AJ, Wallace K, Tsang S, Haile RW, Saibil F, Ahnen D, et al. MTHFR genotype and colorectal adenoma recurrence: data from a double-blind placebo-controlled clinical trial. *Cancer Epidemiol Biomarkers Prev* 2008;17:2409–15.
- Cho E, Zhang X, Townsend MK, Selhub J, Paul L, Rosner B, et al. Unmetabolized folic acid in prediagnostic plasma and the risk of colorectal cancer. *J Natl Cancer Inst* 2015;107:12.
- Ma J, Stampfer MJ, Giovannucci E, Artigas C, Hunter DJ, Fuchs C, et al. Methylenetetrahydrofolate reductase polymorphism, dietary interactions, and risk of colorectal cancer. *Cancer Res* 1997;57:1098–102.
- Kim YI. Folate and colorectal cancer: an evidence-based critical review. *Mol Nutr Food Res* 2007;51:267–92.
- Baggott JE, Vaughn WH, Juliana MM, Eto I, Krumdieck CL, Grubbs CJ. Effects of folate deficiency and supplementation on methylnitrosourea-induced rat mammary tumors. *J Natl Cancer Inst* 1992;84:1740–4.
- Kotsopoulos J, Sohn KJ, Martin R, Choi M, Renslund R, McKerlie C, et al. Dietary folate deficiency suppresses N-methyl-N-nitrosourea-induced mammary tumorigenesis in rats. *Carcinogenesis* 2003;24:937–44.
- Mason JB, Dickstein A, Jacques PF, Haggarty P, Selhub J, Dallal G, et al. A temporal association between folate acid fortification and an increase in colorectal cancer rates may be illuminating important biological principles: a hypothesis. *Cancer Epidemiol Biomarkers Prev* 2007;16:1325–9.
- Ulrich CM, Potter JD. Folate supplementation: too much of a good thing? *Cancer Epidemiol Biomarkers Prev* 2006;15:189–93.
- Wright AJ, Dainty JR, Finglas PM. Folic acid metabolism in human subjects revisited: potential implications for proposed mandatory folic acid fortification in the UK. *Br J Nutr* 2007;98:667–75.
- Obeid R, Herrmann W. The emerging role of unmetabolized folic acid in human diseases: myth or reality? *Curr Drug Metab* 2012;13:1184–95.
- Kelly P, McPartlin J, Goggins M, Weir DG, Scott JM. Unmetabolized folic acid in serum: acute studies in subjects consuming fortified food and supplements. *Am J Clin Nutr* 1997;65:1790–5.
- Morris MS, Jacques PF, Rosenberg IH, Selhub J. Circulating unmetabolized folic acid and 5-methyltetrahydrofolate in relation to anemia, macrocytosis, and cognitive test performance in American seniors. *Am J Clin Nutr* 2010;91:1733–44.
- Hirsch S, Miranda D, Munoz E, Montoya M, Ronco AM, de la Maza MP, et al. Natural killer cell cytotoxicity is not regulated by folic acid in vitro. *Nutrition* 2013;29:772–6.
- Ohrvik VE, Withoft CM. Human folate bioavailability. *Nutrients* 2011;3:475–90.
- Bailey SW, Ayling JE. The extremely slow and variable activity of dihydrofolate reductase in human liver and its implications for high folic acid intake. *Proc Natl Acad Sci U S A* 2009;106:15424–9.
- Christensen KE, Mikael LG, Leung KY, Levesque N, Deng L, Wu Q, et al. High folic acid consumption leads to pseudo-MTHFR deficiency, altered

Rees et al.

- lipid metabolism, and liver injury in mice. *Am J Clin Nutr* 2015;101:646–58.
40. Wright AJ, Finglas PM, Dainty JR, Hart DJ, Wolfe CA, Southon S, et al. Single oral doses of ¹³C forms of pteroylmonoglutamic acid and 5-formyltetrahydrofolic acid elicit differences in short-term kinetics of labelled and unlabelled folates in plasma: potential problems in interpretation of folate bioavailability studies. *Br J Nutr* 2003;90:363–71.
 41. Obeid R, Kasoha M, Kirsch SH, Munz W, Herrmann W. Concentrations of unmetabolized folic acid and primary folate forms in pregnant women at delivery and in umbilical cord blood. *Am J Clin Nutr* 2010;92:1416–22.
 42. Obeid R, Kirsch SH, Kasoha M, Eckert R, Herrmann W. Concentrations of unmetabolized folic acid and primary folate forms in plasma after folic acid treatment in older adults. *Metabolism* 2011;60:673–80.
 43. Kalmbach RD, Choumenkovitch SF, Troen AM, D'Agostino R, Jacques PF, Selhub J. Circulating folic acid in plasma: relation to folic acid fortification. *Am J Clin Nutr* 2008;88:763–8.
 44. Bailey RL, Mills JL, Yetley EA, Gahche JJ, Pfeiffer CM, Dwyer JT, et al. Serum unmetabolized folic acid in a nationally representative sample of adults ≥60 years in the United States, 2001–2002. *Food and Nutrition Research* 2012;56:10.3402/fnr.v56i0.5616.
 45. Centers for Disease Control. Recommendations for the use of folic acid to reduce the number of cases of spina bifida and other neural tube defects. *MMWR* 1992;41(RR-14):1–7.
 46. Brouwer IA, van Dusseldorp M, West CE, Steegers-Theunissen RP. Bioavailability and bioefficacy of folate and folic acid in man. *Nutr Res Rev* 2001;14:267–94.
 47. Alemdaroglu NC, Dietz U, Wolfram S, Spahn-Langguth H, Langguth P. Influence of green and black tea on folic acid pharmacokinetics in healthy volunteers: potential risk of diminished folic acid bioavailability. *Biopharmaceutics Drug Disposition* 2008;29:335–48.
 48. Willems FF, Boers GH, Blom HJ, Aengevaeren WR, Verheugt FW. Pharmacokinetic study on the utilisation of 5-methyltetrahydrofolate and folic acid in patients with coronary artery disease. *British J Pharmacol* 2004;141:825–30.
 49. Loew D, Eberhardt A, Hesecker H, Kubler W. [Plasma kinetics and elimination of folic acid]. *Klin Wochenschr* 1987;65:520–4.
 50. von der Porten AE, Gregory JF III, Toth JP, Cerda JJ, Curry SH, Bailey LB. In vivo folate kinetics during chronic supplementation of human subjects with deuterium-labeled folic acid. *J Nutr* 1992;122:1293–99.
 51. Hutcheon JA, Chioloro A, Hanley JA. Random measurement error and regression dilution bias. *BMJ* 2010;340:c2289.

Cancer Prevention Research

Unmetabolized Folic Acid, Tetrahydrofolate, and Colorectal Adenoma Risk

Judy R. Rees, Carolyn B. Morris, Janet L. Peacock, et al.

Cancer Prev Res 2017;10:451-458. Published OnlineFirst June 9, 2017.

Updated version Access the most recent version of this article at:
doi:[10.1158/1940-6207.CAPR-16-0278](https://doi.org/10.1158/1940-6207.CAPR-16-0278)

Supplementary Material Access the most recent supplemental material at:
<http://cancerpreventionresearch.aacrjournals.org/content/suppl/2017/07/12/1940-6207.CAPR-16-0278.DC1>

Cited articles This article cites 51 articles, 24 of which you can access for free at:
<http://cancerpreventionresearch.aacrjournals.org/content/10/8/451.full#ref-list-1>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.